Functional Ecology 1993
7, 759–762

Author Index

Ackerman J.D. & Okubo A.: Reduced mixing in a marine macrophyte canopy, 305
Adler L.S., Wikler K., Wyndham F.S., Linder C.R. & Schmitt J.: Potential for persistence of genes escaped from canola: germination cues in crop, wild, and crop-wild hybrid *Brassica rapa*, 736
Agarwala B.K. & Dixon A.F.G.: Why do ladybirds lay eggs in clusters? 541
Allen M.F., *see* Friese C.F.
Alm D.M., *see* Nobel P.S.
Asami T.: Interspecific differences in desiccation tolerance of juvenile land snails, 571

Baird D.J.: Can toxicity testing contribute to ecotoxicology? 510
Bale J.S.: Classes of insect cold hardiness, 751
Ballabeni P. & Ward P.I.: Local adaptation of the tremadote *Diplostomum phoxini* to the European minnow *Phoxinus phoxinus*, its second intermediate host, 84
Band S.R., *see* Thompson K.
Baudinette R.V., *see* MacMillen R.E.
Bayne B.L., *see* MacDonald B.A.
Beaupre S.J., Dunham A.E. & Overall K.L.: The effects of consumption rate and temperature on apparent digestibility coefficient, urate production, metabolizable energy coefficient and passage time in canyon lizards (*Sceloporus merriami*) from two populations, 273
Behmer S.T. & Joern A.: Diet choice by a grass-feeding grasshopper based on the need for a limiting nutrient, 522
Bennett A.F., *see* Losos J.B.
Bennett G.F., *see* Korpimäki E.
Berdoy M., *see* Smith P.
Berendse F., *see* Elberse W.Th.
Beyschlag W., *see* Ryel R.J.
Biernacka B., *see* Dratnal E.
Black R.A., *see* Williams D.G.
Bliss L.C., *see* Dawson T.E.
Block W., *see* Wharton D.A.
Blom C.W.P.M., *see* Voesenek L.A.C.J.
Borghetti M., Leonardi S., Raschi A., Snyderman D. & Tognetti R.: Ecotypic variation of xylem embolism, phenological traits, growth parameters and allozyme characteristics in *Fagus sylvatica*, 713
Bowes G., *see* Moon M.
Brinklow B.R., *see* Heydon M.J.
Brown V.K., *see* Gange A.C.
Budaraju S., *see* Stewart W.E.

Cabral O.M.R., *see* McWilliam A.-L.C.
Cagle K.D., Packard G.C., Miller K. & Packard M.J.: Effects of the microclimate in natural nests on development of embryonic painted turtles, *Chrysemys picta*, 653
Caldwell M.M., *see* Ryel R.J.
Chazdon R.L. & Kaufmann S.: Plasticity of leaf anatomy of two rain forest shrubs in relation to photosynthetic light acclimation, 385
Chiba N. & Hirose T.: Nitrogen acquisition and use in three perennials in the early stage of primary succession, 287
Chown S.L.: Desiccation resistance in six sub-Antarctic weevils (Coleoptera: Curculionidae): humidity as an abiotic factor influencing assemblage structure, 318
Clarke A.: Reproductive trade-offs in caridean shrimps, 411
Seasonal acclimatization and latitudinal compensation in metabolism: do they exist? 139
Colasanti R.L. & Grime J.P.: Resource dynamics and vegetation processes: a deterministic model using two-dimensional cellular automata, 169
de Costa A.C.L., *see* McWilliam A.-L.C.
Crawford R.M.M., *see* Studer-Ehrensberger K.

Danckwerts J.E.: Reserve carbon and photosynthesis: their role in regrowth of *Themeda triandra*, a widely distributed subtropical graminaecous species, 634
Davies R.W., *see* Dratnal E.
Dawson T.E. & Bliss L.C.: Plants as mosaics: leaf-, ramet-, and gender-level variation in the physiology of the dwarf willow, *Salix arctica*, 293
Depledge M.H., *see* Forbes V.E.
Dewar R.C.: A root–shoot partitioning model based on carbon–nitrogen–water interactions and Münch phloem flow, 356
Dixon A.F.G., Kundu R. & Kindlmann P.: Reproductive effort and maternal age in iteroparous insects using aphids as a model group, 267
see also Agarwala B.K.
Djawdan M.: Locomotor performance of bipedal and quadrupedal heteromyid rodents, 195
Dratnal E., Reddy D.C., Biernacka B. & Davies R.W.: Facultative physiological adaptation and compensation to winter stresses in the predatory leech *Nephelopsis obscura*, 91
Dufrene E. & Saugier B.: Gas exchange of oil palm in relation to light, vapour pressure deficit, temperature and leaf age, 97
Dumesnil M.J., *see* Mayeux H.S.
Dunham A.E., *see* Beaupre S.J.

Eason W.R., *see* Newman E.I.
Eguiarte L.E., *see* López-Portillo J.
Elberse W.Th. & Berendse F.: A comparative study of the growth and morphology of eight grass species from habitats with different nutrient availabilities, 223
El Kohen A., Venet L. & Mousseau M.: Growth and photosynthesis of two deciduous forest species at elevated carbon dioxide, 480

Forbes T.L. & Forbes V.E.: A critique of the use of distribution-based extrapolation models in ecotoxicology, 249
see also Forbes V.E.
Forbes V.E. & Depledge M.H.: Testing *vs* research in ecotoxicology: a response to Baird and Calow, 509
Forbes V.E. & Forbes T.L.: Ecotoxicology and the power of clones, 511
see also Forbes T.L.
Forsman A. & Lindell L.E.: The advantage of a big head: swallowing performance in adders, *Vipera berus*, 183
Fox C.W.: Multiple mating, lifetime fecundity and female mortality of the bruchid beetle, *Callosobruchus maculatus* (Coleoptera: Bruchidae), 203
Franco M., *see* Silvertown J.
Friese C.F. & Allen M.F.: The interaction of harvester ants and vesicular–arbuscular mycorrhizal fungi in a patchy

semi-arid environment: the effects of mound structure on fungal dispersion and establishment, 13

Gabriel W., *see* Taylor B.E.
Gange A.C., Brown V.K. & Sinclair G.S.: Vesicular–arbuscular mycorrhizal fungi: a determinant of plant community structure in early succession, 616
Giordano P.A. & Hibbs D.E.: Morphological response to competition in red alder: the role of water, 462
Graves J.L., *see* Mueller L.D.
Grime J.P.: A comment on Silvertown, Franco & McConway (1992), 380
An olive branch and further clarification, 381
see also Colasanti R.L.
Grubb P.J., *see* Marañón T.
Guillette L.J., Jr, *see* Méndez-de la Cruz F.R.

Hakkarainen H., *see* Korpimäki E.
Hand D.W., *see* Hunt R.
Hannah M.A., *see* Hunt R.
Hansell M.H.: The ecological impact of animal nests and burrows, 5
Hastings S.J., *see* Vourlitis G.L.
Heydon M.J., Sibbald A.M., Milne J.A., Brinklow B.R. & Loudon A.S.I.: The interaction of food availability and endogenous physiological cycles on the grazing ecology of red deer hinds (*Cervus elaphus*), 216
Hibbs D.E., *see* Giordano P.A.
Hickling G.J. & Millar J.S.: The fasting endurance model: a testable hypothesis, 135
Hirose T., *see* Chiba N.
Hodgson J.G., *see* Thompson K.
Hopkins W.G., *see* Yuan T.
Houston A.I., *see* Weisser W.W.
Houston D.C., *see* Lee W.B.
Hughes L., Westoby M. & Johnson A.D.: Nutrient costs of vertebrate- and ant-dispersed fruits, 54
Hunt R., Hand D.W., Hannah M.A. & Neal A.M.: Further responses to CO_2 enrichment in British herbaceous species, 661

Jaeger J., *see* Stewart W.E.
Jarvis P.G., *see* Wang Y.P.
Jenkins M.A., *see* Vourlitis G.L.
Joern A., *see* Behmer S.T.
Johnson A.D., *see* Hughes L.
Johnson H.B., *see* Mayeux H.S.
Johnston R.D.: Effects of diet quality on the nestling growth of a wild insectivorous passerine, the house martin *Delichon urbica*, 255
Jokela J., Uotila L. & Taskinen J.: Effect of the castrating trematode parasite *Rhipidocotyle fennica* on energy allocation of fresh-water clam *Anodonta piscinalis*, 332
de Jong G.: Covariances between traits deriving from successive allocations of a resource, 75
Judge M.L., *see* Levinton J.S.

Kaufmann S., *see* Chazdon R.L.
Keller L.F. & van Noordwijk A.J.: A method to isolate environmental effects on nestling growth, illustrated with examples from the great tit (*Parus major*), 493
Kindlmann P., *see* Dixon A.F.G.
Koike F. & Syahbuddin: Canopy structure of a tropical rain forest and the nature of an unstratified upper layer, 230
Kolunen H., *see* Pietiäinen H.
Korpimäki E., Hakkarainen H. & Bennett G.F.: Blood parasites and reproductive success of Tengmalm's owls: detrimental effects on females but not on males? 420
Kouki J.: Herbivory modifies the production of different leaf types in the yellow water-lily, *Nuphar lutea* (Nymphaeaceae), 21
Kundu R., *see* Dixon A.F.G.

Lambers H., *see* van der Werf A.
Lee W.B. & Houston D.C.: The role of coprophagy in digestion in voles (*Microtus agrestis* and *Clethrionomys glareolus*), 427
Leitao M.V.B.R., *see* McWilliam A.-L.C.
Leonardi S., *see* Borghetti M.
Lerdau M.: Formal equivalence among resource allocation models: what is the appropriate currency? 507
Levinton J.S. & Judge M.L.: The relationship of closing force to body size for the major claw of *Uca pugnax* (Decapoda: Ocypodidae), 339
Lindell L.E., *see* Forsman A.
Linder C.R., *see* Adler L.S.
Lloyd D.G., *see* Robertson A.W.
López-Portillo J., Eguiarte L.E. & Montaña C.: Nectarless honey mesquites, 452
Losos J.B., Walton B.M. & Bennett A.F.: Trade-offs between sprinting and clinging ability in Kenyan chameleons, 281
Loudon A.S.I., *see* Heydon M.J.
Lovett Doust J., *see* Lovett Doust L.
Lovett Doust L., Lovett Doust J. & Schmidt M.: In praise of plants as biomonitors—send in the clones, 754

McArthur C. & Sanson G.D.: Nutritional effects and costs of a tannin in a grazing and a browsing macropodid marsupial herbivore, 690
McArthur C. & Sanson G.D.: Nutritional effects and costs of a tannin in two marsupial arboreal folivores, 697
McConway K., *see* Silvertown J.
MacDonald B.A. & Bayne B.L.: Food availability and resource allocation in senescent *Placopecten magellanicus*: evidence from field populations, 40
MacDonald D.W., *see* Smith P.
MacGillivray C.W., *see* Thorpe P.C.
Macko S.A., *see* Wallace L.L.
MacMillen R.E. & Baudinette R.V.: Water economy of granivorous birds: Australian parrots, 704
McNeil J.N., *see* Royer L.
McWilliam A.-L.C., Roberts J.M., Cabral O.M.R., Leitao M.V.B.R., de Costa A.C.L., Maitelli G.T. & Zamparoni C.A.G.P.: Lead area index and above-ground biomass of *terra firme* rain forest and adjacent clearings in Amazonia, 310
Maitelli G.T., *see* McWilliam A.-L.C.
Marañón T. & Grubb P.J.: Physiological basis and ecological significance of the seed size and relative growth rate relationship in Mediterranean annuals, 591
Maun M.A., *see* Yuan T.
Mayeux H.S., Johnson H.B., Polley H.W., Dumesnil M.J. & Spanel G.A.: A controlled environment chamber for growing plants across a subambient CO_2 gradient, 125
Méndez-de la Cruz F.R., Guillette L.J., Jr & Villagran-Santa Cruz M.: Differential atresia of ovarian follicles and its effect on the clutch size of two populations of the viviparous lizard *Sceloporus mucronatus*, 535

Menozzi P., *see* Rossi V.
Millar J.S., *see* Hickling G.J.
Miller K., *see* Cagle K.D.
Milne J.A., *see* Heydon M.J.
Montaña C., *see* López-Portillo J.
Moon M., Rattray M.R., Putz F.E. & Bowes G.: Acclimatization to flooding of the herbaceous vine, *Mikania scandens*, 610
Mousseau M., *see* El Kohen A.
Mueller L.D., Graves J.L. & Rose M.R.: Interactions between density-dependent and age-specific selection in *Drosophila melanogaster*, 469

Neal A.M., *see* Hunt R.
Newman E.I. & Eason W.R.: Rates of phosphorus transfer within and between ryegrass (*Lolium perenne*) plants, 242
Nobel P.S. & Alm D.M.: Root orientation *vs* water uptake simulated for monocotyledonous and dicotyledonous desert succulents by a root-segment model, 600
van Noordwijk A.J., *see* Keller L.F.

Obeso J.R.: Does defoliation affect reproductive output in herbaceous perennials and woody plants in different ways? 150
Oechel W.C., *see* Vourlitis G.L.
Okubo A., *see* Ackerman J.D.
van Oorschot F.J.M.M., *see* Voesenek L.A.C.J.
Orozco-Segovia A., Sanchez-Coronado M.E. & Vazquez-Yanes C.: Effect of maternal light environment on seed germination in *Piper auritum*, 395
Light environment and phytochrome-controlled germination in *Piper auritum*, 585
Overall K.L., *see* Beaupre S.J.

Packard G.C., *see* Cagle K.D.
Packard M.J., *see* Cagle K.D.
Pearcy R.W., *see* Roden J.S.
see also Sims D.A.
Perrin N.: On future discounts and economic analogy in life-history studies, 506
Pietiäinen H. & Kolunen H.: Female body condition and breeding of the Ural owl *Strix uralensis*, 726
Polley H.W., *see* Mayeux H.S.
Porter W.P., *see* Stewart W.E.
Priestman G.H., *see* Thorpe P.C.
Purvis A., *see* Stockley P.
Putz F.E., *see* Moon M.

Raschi A., *see* Borghetti M.
Rattray M.R., *see* Moon M.
Reddy D.C., *see* Dratnal E.
Reich P.B.: Reconciling apparent discrepancies among studies relating life span, structure and function of leaves in contrasting plant life forms and climates: 'the blind men and the elephant retold', 721
Roberts J.M., *see* McWilliam A.-L.C.
Robertson A.W. & Lloyd D.G.: Rates of pollen deposition and removal in *Myosotis colensoi*, 549
Roden J.S. & Pearcy R.W.: The effect of leaf flutter on the flux of CO_2 in poplar leaves, 669
Rose M.R., *see* Mueller L.D.
Rossi V. & Menozzi P.: The clonal ecology of *Heterocypris incongruens* (Ostracoda): life-history traits and photo period, 177
Rowell-Rahier M., *see* Speiser B.
Royer L. & McNeil J.N.: Male investment in the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae): impact on female longevity and reproductive performance, 209
Ryel R.J., Beyschlag W. & Caldwell M.M.: Foliage orientation and carbon gain in two tussock grasses as assessed with a new whole-plant gas-exchange model, 115

Salamolard M. & Weimerskirsch H.: Relationship between foraging effort and energy requirement throughout the breeding season in the wandering albatross, 643
Sanchez-Coronado M.E., *see* Orozco-Segovia A.
Sanson G.D., *see* McArthur C.
Saugier B., *see* Dufrene E.
Schieving F., *see* van der Werf A.
Schmidt M., *see* Lovett Doust L.
Schmitt J., *see* Adler L.S.
Sibbald A.M., *see* Heydon M.J.
Silvertown J., Franco M. & McConway K.: The eternal triangle — an attempt at reconciliation, 380
Sims D.A. & Pearcy R.W.: Sunfleck frequency and duration affects growth rate of the understorey plant, *Alocasia macrorrhiza*, 683
Sinclair G.S., *see* Gange A.C.
Smith P., Berdoy M., Smith R.H. & MacDonald D.W.: A new aspect of warfarin resistance in wild rats: benefits in the absence of poison, 190
Smith R.H., *see* Smith P.
Smits A.J.M., *see* Voesenek L.A.C.J.
Snyderman D., *see* Borghetti M.
Spanel G.A., *see* Mayeux H.S.
Speakman J.R.: Evolution of body size; predictions from energetics, 134
How should we calculate CO_2 production in doubly labelled water studies of animals? 746
Speiser B. & Rowell-Rahier M.: Does the land snail *Arianta arbustorum* prefer sequentially mixed over pure diets? 403
Stewart W.E., Budaraju S., Porter W.P. & Jaeger J.: Prediction of forced ventilation in animal fur under ideal pressure distribution, 487
Stockley P. & Purvis A.: Sperm competition in mammals: a comparative study of male roles and relative investment in sperm production, 560
Studer C., *see* Studer-Ehrensberger K.
Studer-Ehrensberger K., Studer C. & Crawford R.M.M.: Competition at community boundaries: mechanisms of vegetation structure in a dune–slack complex, 156
Syahbuddin, *see* Koike F.

Taskinen J., *see* Jokela J.
Taylor B.E. & Gabriel W.: Optimal adult growth of *Daphnia* in a seasonal environment, 513
Thompson K., Band S.R. & Hodgson J.G.: Seed size and shape predict persistance in soil, 236
Thorpe P.C., MacGillivray C.W. & Priestman G.H.: A portable device for the simulation of air frosts at remote field locations, 503
Tognetti R., *see* Borghetti M.

Uotila L., *see* Jokela J.

Vazquez-Yanes C., *see* Orozco-Segovia A.
Venet L., *see* El Kohen A.

Villagran-Santa Cruz M., *see* Méndez-de la Cruz F.R.
Visser A.J., *see* van der Werf A.
Voesenek L.A.C.J., van Oorschot F.J.M.M., Smits A.J.M. & Blom C.W.P.M.: The role of flooding resistance in the establishment of *Rumex* seedlings in river flood plains, 105
Vourlitis G.L., Oechel W.C., Hastings S.J. & Jenkins M.A.: A system for measuring *in situ* CO_2 and CH_4 flux in unmanaged ecosystems: an arctic example, 369

Wallace L.L. & Macko S.A.: Nutrient acquisition by clipped plants as a measure of competitive success: the effects of compensation, 326
Walton B.M., *see* Losos J.B.
Wang Y.P. & Jarvis P.G.: Influence of shoot structure on the photosynthesis of Sitka spruce (*Picea sitchensis*), 433
Ward P.I., *see* Ballabeni P.
Wardle D.A.: Changes in the microbial biomass and metabolic quotient during leaf litter succession in some New Zealand forest and scrubland ecosystems, 346
Weimerskirsch H., *see* Salamolard M.
Weiner J.: Competition, herbivory and plant size variability: *Hypochaeris radicata* grazed by snails (*Helix aspersa*), 47
Weisser W.W. & Houston A.I.: Host discrimination in parasitic wasps: when is it advantageous? 27
van der Werf A., Visser A.J., Schieving F. & Lambers H.: Evidence for optimal partitioning of biomass and nitrogen at a range of nitrogen availabilities for a fast- and slow-growing species, 63
Westoby M., *see* Hughes L.
Wharton D.A. & Block W.: Freezing tolerance in some Antarctic nematodes, 578
Wikler K., *see* Adler L.S.
Williams D.G. & Black R.A.: Phenotypic variation in contrasting temperature environments: growth and photosynthesis in *Pennisetum setaceum* from different altitudes on Hawaii, 623
Wyndham F.S., *see* Adler L.S.

Yuan T., Maun M.A. & Hopkins W.G.: Effects of sand accretion on photosynthesis, leaf-water potential and morphology of two dune grasses, 676

Zamparoni C.A.G.P., *see* McWilliam A.-L.C.

Functional Ecology 1993
7, 763–767

Subject Index

Adalia bipunctata 541–8
ADC *see* apparent digestibility coefficient
adder 183–9, *see also* reptiles
Adenostyles alliariae 403–9
adult density, in fruit flies 469–78
Aegolius funereus 420–5
aerenchyma 610–14
Agave deserti 600–8
ageing, in fruit flies 469–78
Agropyron desertorum 115–23
albatross, wandering 643–52
alder, red, competition and water 462–8
allocation 63–73, 75–83, 223–8, 356–68
allometry, intraspecific 560–8
allozyme characteristics, in beech 713–20
Alnus rubra 462–8
Alocasia macrorrhiza, and sunflecks 683–9
Alopecurus pratensis 223–8
altitudinal gradient 623–31
Amazonia 310–16
amino acid, aromatic 522
Ammophila breviligulata 676–82
analysis of variance 230–4
Andropogon gerardii 326–31
animal fur, forced ventilation in 487–92
animal nests and burrows 5–10
animal–plant interactions 47–52
animals, and carbon dioxide production 746–50
annuals 591–8, *see also* perennials
Anodonta piscinalis 332–7
anoxia tolerance 156–67
Antarctonesiotes elongatus 318–24
Anthoxanthum odoratum 223–8
ants 13–19, 54–9
aphids 267–71
Apis mellifera 452–60
apparent digestibility coefficient (ADC) 273–9
Arianta arbustorum 403–9
arils 54–9
Arrhenatherum elatius 223–8
Artemisia tridentata 13–19
aspen, leaf flutter of 669–75, *see also* trees
Aster ageratoides 287–91
ATP generation 142
atresia, follicular 535–9
automimicry 452–60

beech 480–5, 713–20, *see also* trees
bees 452–60
beetles 203–7, *see also* insects
behaviour, in red deer 216–20
biomass 310–16, 346–54
biomass allocation 623–31
biomechanics 305–8, 339–44
biomes 721–5
birds, granivorous, water and 704–12
body size, in nestlings 493–501
body water 318–24, 571–7
body weight, of rats 190–4
boundaries 156–67
boundary layer 669–75
Brassica rapa, germination in 736–45
Briza media 63–73
bulk flow 669–75
bundle sheath cells 676

Calamovilfa longifolia 676–82
Callosobruchus maculatus 203–7
canopy 305–8, 385–94, 433–51
carbon allocation pattern 682–9
carbon assimilation 293
carbon balance 125–32
carbon gain, in tussock grasses 115–23
carbon and nitrogen substrates 356–68
carbon reserves, in *Themeda triandra* 634–40
carbon–nitrogen–water interactions 356–68
Carex arenaria 156–67
Carex nigra 156–67
Castanea sativa 480–5
Cavariella aegopodii 267–71
Cervus elaphus 216–20
Chamaeleo dilepis 281–6
Chamaeleo jacksonii 281–6
chameleons, Kenyan 281–6
chestnut, sweet 480–5, *see also* trees
Chorismus antarcticus 411–17
Chrysemys picta 653–60
Circaea lutetiana 403–9
clams, freshwater 332–7
Clethrionomys glareolus 427–31
climate 369–77, 503–5, 549–58, 661–8
clipping 326–31, 634–40
clones 177–82, 754–8
clutch size 420–5, 535–9, 726–35
CO_2 assimilation 433–512
CO_2 enrichment 661–8
CO_2 exchange 669–75
CO_2 fixation 683–9
CO_2 gradient, subambient 125–32
CO_2 production 369–77, 480–5, 746–50
co-limitation 433–51
Coccinella septempunctata 541–8
coevolution 84–9
communities 156–67
community organization 318–24
community response 503–5
compensation, and competition 326–31
competition 47–52, 125–32, 156–67, 326–31
conifers 433–51, *see also* trees
convection, forced 487–92
Coomansus gerlachei 578–84
coprophagy, in voles 427–31
covariances between traits 75–83
Crustacea 339–44, 411–17
cryomicroscopy 578–84
CSR theory 169–76

Dactylis glomerata 63–73
Daphnia sp., optimal adult growth of 513–20
Darcy's law 487–92

deer, red 216–20, 490
defoliation 150–3, 634–40
deforestation 310–16
Delichon urbica 255–64
density dependence, seed germination 736–45
desiccation resistance, in weevils 318–24
diaspore dimension 236–40
dicotyledons, succulent 600–8
diet
  of grass-feeding grasshopper 522–6, *see also* food
  of nestling house martins 255–64
  sequentially mixed and pure 403–9
digestion 273–9
  in marsupials 690–6, 697–703
Diplostomum phoxini 84–9
dispersal spectra 54–9
disturbance 13–19, 169–76
Ditylenchus sp. 578–84
docks 105–12
Drosophila melanogaster 469–78
dry matter 310
dry weight partitioning 480–5
dune–slack complex 156–67
dynamic programming 27–37

ecology, clonal 177–82
ecotoxicology
  models in 249–53
  and power of clones 511–12
  testing *vs* research 509–10
  and toxicology testing 510–11
ecotypes 713–20
Ectemnorhinus group 318–24
eddy viscosity 305–8
eelgrass canopy 305–8
egg clustering, of ladybirds 541–8
eggs
  of painted turtles 653–60
  in shrimps 411–17
ejaculates 290–14, 560–8
Elaeis guineensis 97–102
elaiosomes 54–9
electrophoresis 177–82
energetics
  ecological, and digestion 273–9
  evolution of body size 134
energy acquisition 91–6
energy allocation 40–5, 91–6, 267–71
energy coefficient, metabolizable 273–9
energy requirement, of albatross 643–52
energy storage, in clams 332–7
environmental variation, and nestlings 493–501
ESS *see* evolutionarily stable strategy
ethanolic fermentation 105–12
etiolated regrowth 634–40
European corn borer 209–14
evolutionarily stable strategy (ESS) 560–8
evolutionary physiology, in fruit flies 469–78
exponential growth equations 361–2

facultative response 91–6
Fagus sylvatica 480–5, 713–20
fasting endurance model 135
fecundity 411–17
feeding behaviour, in snails 403–9
Ferocactus acanthodes 600–8
Festuca arundinacea 223–8
Festuca ovina 223–8
Festuca protensis 223–8
Festuca rubra 223–8
fiddler crab claw, closing force of 339–44
field establishment 105–12
field metabolic rate 255–64
field studies 503–5
Filipendula ulmaria 156–67
First Major Allocation 75–83
fish 84–9
fitness 513–20, 528
fledgling weight 493–501
flood resistance, and herbaceous vines 610–14
flooding 156–67
fly, tachinid 549, *see also* insects
foliage 115–23, 310–16
  density of 230–4
folivores, marsupial 697–703
food 7, 40–5, 216–20, *see also* diet
food availability 40–5, 216–20
foraging efforts, of wandering albatross 643–52
foundation theory 10
freeze avoidance, in nematodes 578–84
frugivory 54–9, *see also* herbivory
fruit flies 469–78
fruits 54–9
functional balance 356–68
functional types 661–8
fungi, vesicular–arbuscular mycorrhizal (VAM) 13–19, 616–21

Galerucella nymphaeae 21–5
gape limitation 183–9
gas exchange 115–23, 634–40
gastropod 47–52, 403–9, 571–7
gender-level variation, in willow 293–308
gene flow, in *Brassica rapa* 736–45
genetic variation, and nestlings 493–501
genetics 84–9
genets 623–31
genotypic correlations 528–33
germination 585–90, 736–45
Gigaspora margarita 13–19
global change 125–32, 369–77
Glyceria maxima 156–67
gradients, of vegetative disturbance 169–76
grass species, growth and morphology of 223–8
grasses 115–23, 242–7, 326–31, 676–82
grasshopper, diet choice by 522–6
grasslands 326–31
growth 145–6
  adult, of *Daphnia* sp. 513–20
  in deciduous forest species 480–5
  in *Pennisetum setaceum* 623–31
  root and shoot 361
growth analysis 287–91
  in red alder 462–8
growth components, of annuals 591–8
growth equations 493–501

habitat saturation theory 10

habitats
  alteration of 5–7
  dry, rockface 318–24
  invasion of new 8–9
  nutrient-rich and poor 223–8
Haemoproteus noctuae 420–5
Haemoproteus syrnii 420–5
Hamilton and Zuk hypothesis 84–9, 424–5
heat generation 142
Helix aspersa 47–52
herbaceous species, CO_2 enrichment in 661–8
herbivore
  graminivorous 522–6
  marsupial, and tannin 690–6
herbivores 634–40
  generalist 403–9
herbivory 21–5, 47–52, 326–31, 427–31, *see also* frugivory
Heterocypris incongruens, clonal ecology of 177–82
heterogeneity 395–401
heterophylly 21–5
honey mesquites, nectarless 452–60
host reproduction 332–7
host–parasite interaction 84–9
house martin 255–64, *see also* birds
hydraulic conductivity, in succulents 600–8
hydroelasticity 305–8
hyperbolic response 661–8
Hypochaeris radicata 47–52
hypoxia 91–6, 105–12

ice nucleation, exogenous 578–84
in situ measurements 369–77
Indian meal moth 528–33
insects 267–71, 751–3
integration, phenotypic 75–83
invertebrates, benthic suspension feeding 40
iprodione 616–21
isoenzymes 713–20

kangaroo mouse 195
kangaroo rat 195–202

ladybirds, clustering of eggs 541–8
leaf, hypostomatous 669–75
leaf age 293–308
leaf anatomy, plasticity of 385–94
leaf area 310–16, 721–5
leaf flutter, and CO_2 flux 669–75
leaf life span 721–5
leaf litter 346–54
leaf nitrogen 721–5
leaf structure 385–94
leaf thickness 676–82
leaf variation 293–308
leaf-to-air vapour pressure differences 293
leaf-water potential 676–82
leucocytozoids 420–5
Leucocytozoon ziemanni 420–5
life history
  of *Daphnia* sp. 513–20
  in shrimps 411–17
life-history evolution 469–78
life-history studies 506–7
life-history theory 267–71
life span, of bruchid beetles 203–7
light, uniform diffuse 683–9
light climate 683–9
lizards 273–9, 535–9
Lolium perenne 223–8, 242–7

macroclimate 346–54
Macropus parma 690–6
malatonin 216–20
mammals 487–92, 560–8, *see also* animals
marsupials 690–6, 697–703
maternal age, in iteroparous insects 267–71
maternal effects 736–45
mating 203–7, 420–5
MEC *see* metabolizable energy coefficient
Megoura viciae 267–71
Melissodes tristis 452–60
Mesodon normalis, desiccation of 571–7
mesquites, nectarless honey 452–60, *see also* trees
metabolic acclimatization 139–48
metabolizable energy coefficient (MEC) 273–9
microflora 346–54
Microtus agrestis 427–31
Mikania scandens 610–14
mineral deficit 365–6
mineral nutrition 223–8
minnow, European 84–9
Miscanthus oligostachyus 287–91
monocotyledons, succulent 600–8
mortality risks 27–37
multiparasitism 27–37, *see also* superparasitism
Münch flow hypothesis 356–68
mutualism 13–19
mycorrhizae 13–19, 242–7, 616–21
Myosotis colensoi 549–58
myrmecochory 54–9, *see also* ants

neighbourhoods 326–31
Nelder density series 462–8
Nematocarcinus lanceopes 411–17
nematodes, Antarctic 578–84
Nephelopsis obscura 91–6
nest material 7
nestlings 255–64, 493–501
nests 5–10, 653–60
net assimilation rate 462–8
New Zealand forest 346–54
nitrogen 54–9
  and marsupials 690–6, 697–703
  and perennials 287–91
nitrogen productivity (NP) 63–73, 287–91
nitrogen uptake 326–31
normoxia 91–6
Notocrangon antarcticus 411–17
NP *see* nitrogen productivity
Nuphar lutea 21–5
nutrient acquisition, by clipped plants 326–31
nutrient transfer, in ryegrass 242–7
nutrients, ejaculate-derived 203–7
nutrition, in nestlings 493–501

obligate parthenogenesis 177–82
Odum's theory of ecosystem succession 169, 346–54
offspring size, of aphids 267–71

oil palm, gas exchange of 97–102
Oklahoma, prairies of 326–31
optimization, of biomass and nitrogen partitioning 63–73
orthoptera *see* grasshoppers
Oryzopsis hymenoides 13–19
Ostrinia nubilalis 209–14
oviposition 27–37, 541–8
owls 420–5, 726–35, *see also* birds
oxygen consumption 141–2
oxygen demand, seasonal changes in 142–3

Palirhoeus eatoni 318–24
Panagrolaimus sp. 578–84
parasites 332–7, 420–5
parasitoids, solitary 27–37
parrots 704–12, *see also* birds
Parus major 493–501, *see also* birds
pathogens, resistance to 528–33
Pennisetum setaceum 623–31
perennials 150–3, 287–91, 616–21, *see also* annuals
permeability, in animal fur 487–92
permeability coefficients 669–75
PFD *see* photon flux density
phagostimulant 522–6
phenology 177–82, 549–58, 713–20
phenylalanine, and grasshoppers 522
Phleum pratense 326–31
Phoetaliotes nebrascensis 522–6
photon flux density (PFD) 386–94
photosynthesis 97–102, 125–32, 223–8, 293–308, 385–94, 623–31, 721–5
  in deciduous forest species 480–5
  and dune grasses 676–82
  Li-Cor portable system 369–77
  and reserve carbon 634–40
  in Sitka spruce 433–51
  and sunflecks 683–9
  underwater 105–12
photosynthesis model, plant 115–23
photosynthetic induction state 683–9
photosynthetically active radiation (PAR) 97–102
Phoxinus phoxinus 84–9
phytochrome 395–401
Picea sitchensis 433–51, *see also* trees
Piper arieianum 385–94
Piper auritum 395–401, 585–90
Piper sancti-felicis 385–94
Placopecten magellanicus 40–5
plant architecture 115–23
plant growth. herbaceous 661–8
plant nitrogen concentration 287–91
plant resource allocation 506–7
plant species richness 616–21
plant strategies 223–8, 661–8
plant–animal interactions 21–5, 47–52
plants
  clipped, nutrient acquisition by 326–31, 634–40
  woody, and defoliation 150–3
plants as biomonitors 754–8
Plasmodium circumflexum 420–5
plasticity, phenotypic 91–6, 623–31
Plodia interpunctella 528–33
PNC *see* plant nitrogen concentration
Pogonomyrmex occidentalis 13–19
poison, and rats 190–4
pollen presentation 549–58
pollination, insect, in honey mesquite 452–60
Polygonum cuspidatum 287–91
poplar leaves, leaf flutter of 669–75
Populus fremontii 669–75, *see also* trees
Populus tremuloides 669–75
porous media 487–92
possum, brushtail 490
potassium 54–9
predation, non-selective 513–20
predator–prey size relationship 183–9
primary succession 287–91
Prosopis glandulosa var. *torreyana* 452–60
Pseudocheirus peregrinus 697–703
Pseudoroegneria spicata 115–23
Pyrrhalta sp. 21–5

quantum flux, incident 433–51
quebracho 690
quotient 736–45

r and K selection 469–78
rain forest 230–4, 310–16, 385–90, 585–90
ramets 293–308
rate maximizing 27–37
Rattus norvegicus 190–4, *see also* rodents
red:far-red (R:FR) 585–90, 736–45
relative growth rate (RGR) 63–73, 287–91, 462–8, 591–8, 610–14
reproduction 40–5, 216–20, 267–71
reproductive success, in European corn borer 209–14, 726–35
reptiles 183–9, 273–9, 535–9
resource allocation 40–5, 75–83, 507–8, 513–20
resource dynamics 169–76
respiration 141, 144–5, 293, 369–77
  anaerobic 610–14
  dark 293–308, 385–94
RGR *see* relative growth rate
Rhipidocotyle fennica 332–7
Richards growth curve 493–501
rodents 190–4, 195–202
root mats 13–19
root orientation 600–8
root-segment model 600–8
root–shoot partitioning model 356–68
Rumex acetosa 105–12
Rumex crispus 105–12
Rumex palustris 105–12
running endurance, of heteromyid rodents 195–202

Salix arctica 293–308
sand accretion, and dune grasses 676–82
satellite telemetry 643–52
scallop, sea 40–5
Sceloporus merriami 273–9
Sceloporus mucronatus 535–9
seagrass 305–8
seasonality, in red deer 216–20
seed bank 236–40
seed germination 395–401, 585–90
seed hydration 395–401
seed maturation 395–401

seed size and shape 236–40
Senecio fuchsii 403–9
sex-specific differences, in albatross 643–52
sexual selection 209–14, 339–44
shade 683–9
shells, snail 571–7
shoot structure, in Sitka spruce 433–51
shrimps, polar caridean 411–17
shrubs, rain forest 385–94
Silvertown, Franco & McConway paper, comments 380–2
Sitka spruce 433–51
SLA *see* specific leaf area
snails 47–52, 403–9, 571–7
snakes 183–9, *see also* reptiles
soil, seed persistence in 236–40
soil fertility 54–9
soil water potential 600–8
species 318–24
specific leaf area (SLA) 223–8, 591–8
specific root length (SRL) 223–8, 287–91
spectral analysis 305–8
sperm competition, in mammals 560–8
spermatophore 209–14
sprint speed, in chameleons 281–6
SRL *see* specific root length
Stachys silvatica 403–9
stomata 97–102, 293–308, 610–14
stratification, of rain forest canopy 230–4
stress 91–6, 252
stress tolerance 156–67
Strix uralensis 726–35
substrates 356–68
succession 169–76, 616–21
succulents 600–8
sunflecks 585–90, 683–9
superparasitism 27–37
survivorship, of beetles 203–7
swallowing capacity, in adders 183–9

tannin, and marsupials 690–6, 697–703
temperature 503–5, 653–60
  ambient 704
  influence of 147–8
  and metabolic rate 143–4
temperature environments 623–31
Tengmalm's owls 420–5
Teratocephalus tilbrooki 578–84
terra firme rain forest 310–16
testis size, in mammals 560–8
Themeda triandra 634–40
Thylogale billardierii 690–6
tissue water relations 293–308
tit, great 493–501, *see also* birds
TNC *see* total non-structural carbohydrate
total non-structural carbohydrate (TNC) 293
trade-offs 40–5, 75–83
  in chameleons 281–6
  reproductive, in shrimps 411–17
  with resistance to pathogens 528–33
transpiration 97–102, 115–23
trematodes, parasitic 84–9, 332–7
Trichosurus vulpecula 697–703
Triodopsis albolabris 571–7
tropical rain forest 230–4, 385–94, 395–401, 585–90
tundra 369–77
turbulence 305–8
turtles, painted, 653–60, *see also* reptiles

Uca pugnax 339–44
ULR *see* unit leaf rate
Unionidae 332–7
unit leaf rate 591–8
urate production 273–9

VAM *see* vesicular–arbuscular mycorrhizas
variable season 513–20
variation 47–52, 293–308
vegetation disturbance, gradients of 169–76
vegetation succession 346–54
velocity profiles 487–92
vesicular–arbuscular mycorrhizal (VAM) 13–19, 616–21
vine, herbaceous 610–14
Vipera berus 183–9
virus, granulosis 538–33
voles 427–31, 726–35
vortex tubes 503–5

warfarin and rats 190–4
wasps 27–37, *see also* insects
water
  doubly labelled 255–64, 746–50
  and red alder 462–8
water balance 318–24, 360–1, 571–9
water potential, and nests of turtles 653–60
water regulatory efficiency 704–12
water uptake, simulated, and root orientation 600–8
water-use efficiency 115–23, 125–32
weevils, sub-Antarctic 318–24
willow, dwarf 293–308, *see also* trees

xylem, in succulents 600–8
xylem embolism, in beech 713–20

Yellowstone National Park 326–31
yolk size, in shrimps 411–17

Zostera marina 305–8